

V 型潘式纳米管线图的特定 Banhatti 指数

瓦格斯·纳泽尔

(拉合尔教育大学 科学与技术部, 拉合尔 54000)

摘要: 拓扑指数是与图表关联的数字量, 其表征图的拓扑在图自同构下是不变的. 拓扑指数被用作结构 - 活性定量关系 (QSAR) 的例子, 其中分子的生物活性或其他性质与其化学结构相关. 因此, 计算了 V 型潘氏纳米管线图的第 1 和第 2 的 K Banhatti 指数, 以及 V 型潘氏纳米管线图的第 1 和第 2 的 K 超 Banhatti 指数.

关键词: K Banhatti 指数; K 超 Banhatti 指数; 线图; 亚苯基; V 型潘式纳米管

中图分类号: O157.6 **文献标识码:** A **文章编号:** 1674 - 5639 (2018) 03 - 0042 - 05

DOI: 10.14091/j.cnki.kmxyxb.2018.03.008

The Certain Banhatti Indices of the Line Graph of V-pantacenic Nanotube

WAQAS Nazeer

(Division of Science and Technology, University of Education Lahore, Lahore, Pakistan 54000)

Abstract: A topological index is a numeric quantity associated with a graph which characterizes the topology of the graph and is invariant under graph automorphism. Topological indices are used as an example in the development of quantitative structure-activity relationships (QSARs) in which the biological activity or other properties of molecules are correlated with their chemical structure. The aim of this report is to compute the first and second K Banhatti indices of the Line Graphs of V-Pantacenic Nanotubes. We also compute the first and second K hyper Banhatti indices of the Line Graphs of V-Pantacenic Nanotubes.

Key words: K Banhatti index; K-hyper Banhatti index; line graph; phenylenes; V-Pantacenic

0 Introduction

Chemical graph theory is a branch of graph theory in which a chemical compound is represented by a simple graph called molecular graph in which vertices are atoms of compound and edges are the atomic bounds. A graph is connected if there is at least one connection between its vertices. Throughout this paper we take G a connected graph. If a graph does not contain any loop or multiple edges then it is called a network. Between two vertices u and v , the distance is the shortest path between them and is denoted by $d(u, v) = d_G(u, v)$ in graph G . For a vertex v of G the “degree” d_v is the number of vertices attached to it. The edge connecting the vertices u and v will be denoted by uv . Let $d_G(e)$ denote the degree of an edge e in G , which is defined by $d_G(e) = d_G(u) + d_G(v) - 2$ with $e = uv$. The degree and valence in chemistry are closely related to each other. We refer to the book^[1] for more details. Another emerging field is Cheminformatics, which helps to predict biological activities with the relationship of Structure-property and quantitative structure-activity. Topological indices and Physico-chemical properties are used in prediction of bioactivity if underlined compounds are used in these studies^[2-3].

The number that describes the topology of a graph is called topological index. In 1947, the first and most studied topological index was introduced by Wiener^[4]. More details about this index can be found in^[5-6]. In 1975, Milan Randic introduced the Randic index^[7].

收稿日期: 2018 - 02 - 23

作者简介: 瓦格斯·纳泽尔, 男, 巴基斯坦人, 副教授, 博士, 主要从事图论和理论化学研究.

Bollobas et al.^[8] and Amic et al.^[9] in 1998, working independently defined the generalized Randić index. This index was studied by both mathematicians and chemists^[10]. For details about topological indices, we refer to^[11–15]. The first and second K-Banhatti indices of G are defined as

$$B_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)],$$

$$B_2(G) = \sum_{uv \in E(G)} [d_G(u) \times d_G(v)],$$

where uv means that the vertex u and edge e are incident in G . The first and second K-hyper Banhatti indices of G are defined as

$$HB_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2,$$

$$HB_2(G) = \sum_{uv \in E(G)} [d_G(u) \times d_G(v)]^2.$$

We refer to^[16–17] for details about these indices.

The first and second multiplicative K Banhatti indices are defined as^[18]

$$BII_1(G) = \prod_{uv} [d_G(u) + d_G(v)],$$

$$BII_2(G) = \prod_{uv} [d_G(u) d_G(v)].$$

The first and second multiplicative K hyper Banhatti indices are defined as^[18]

$$HBII_1(G) = \prod_{uv} (d_G(u) + d_G(v))^2,$$

$$HBII_2(G) = \prod_{uv} (d_G(u) d_G(v))^2.$$

The line graph $L(G)$ of a graph G is the graph each of whose vertices, representing an edge of G and two of its vertices are adjacent if their corresponding edges are adjacent in G .

Lemma 1 Let G be a graph of order p and size q . Then the line graph $L(G)$ of G is a graph of order p and size $\frac{1}{2}M_1(G) - q$.

In this report we compute Banhatti indices of line graph of V-Pantacenic nanotube^[19]. The graph of V-Pantacenic nanotube is given in figure 1 and the line graph of V-Pantacenic nanotube is given in figure 2.




Figure 1 The V-Pantacenic nanotube $F[2,5]$




Figure 2 The line graph of V-Pantacenic nanotube $F[2,5]$

1 Computational Results

In this section we will give our main results.

Theorem 1 Let $G = L(F[p, q])$ be a line graph of V-Pantacenic nanotube. Then

$$B_1(G) = 1\,320pq - 568p,$$

$$B_2(G) = 3\,168pq - 1\,652p,$$

$$HB_1(G) = 13\,200pq - 6\,936p,$$

$$HB_2(G) = 76\,032pq - 49\,132p.$$

Proof The line graph of V-Pantacenic Nanotubes is shown in figure 2. It can be observed from the figure 2 and lemma 1 that

$$|V(G)| = 33pq - 5p,$$

There are two types of vertices based on the degree, i. e

$$V_1(G) = \{v \in V(G) : d_v = 3\},$$

$$V_2(G) = \{v \in V(G) : d_v = 4\},$$

Such that

$$|V_1(G)| = 20p,$$

$$|V_2(G)| = 33pq - 25p.$$

Also

$$|E(G)| = 66pq - 20p.$$

We can divide the edge set of the line graph of V-Pantacenic Nanotube into following three classes depending on the degree of end vertices of each edge;

$$E_{\{3,3\}}(G) = \{e = uv \in E(G) : d_u = 3, d_v = 3\},$$

$$E_{\{3,4\}}(G) = \{e = uv \in E(G) : d_u = 3, d_v = 4\},$$

and

$$E_{\{4,4\}}(G) = \{e = uv \in E(G) : d_u = 4, d_v = 4\}.$$

Now

$$|E_{\{3,3\}}(G)| = 18p,$$

$$|E_{\{3,4\}}(G)| = 20p,$$

$$|E_{\{4,4\}}(G)| = 66pq - 58p.$$

The detailed edge set dividing can refer to Table 1.

Tab. 1 The edge set dividing of $L(F[p, q])$

$d_{(G)}(u), d_{(G)}(v) = e \in E(G)$	$d_{(G)}(e)$	Numbers of edges
(3, 3)	4	18p
(3, 4)	5	20p
(4, 4)	6	66pq - 58p

1) From the definition of first K Banhatti index

$$B_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)] = \sum_{ue \in E_{\{3,3\}}} [(d_G(u) + d_G(e)) + (d_G(v) + d_G(e))] + \sum_{ue \in E_{\{3,4\}}} [(d_G(u) + d_G(e)) + (d_G(v) + d_G(e))] + \sum_{ue \in E_{\{4,4\}}} [(d_G(u) + d_G(e)) + (d_G(v) + d_G(e))] = 18p[(3+4) + (3+4)] + 20p[(3+5) + (4+5)] + (66pq - 58p)[(4+6) + (4+6)] = 1320pq - 568p.$$

2) From the definition of second K Banhatti index

$$B_2(G) = \sum_{uv \in E(G)} [d_G(u) \times d_G(v)] = \sum_{ue \in E_{\{3,3\}}} [(d_G(u)d_G(e)) + (d_G(v)d_G(e))] + \sum_{ue \in E_{\{3,4\}}} [(d_G(u)d_G(e)) + (d_G(v)d_G(e))] + \sum_{ue \in E_{\{4,4\}}} [(d_G(u)d_G(e)) + (d_G(v)d_G(e))] = 12[(3 \times 4) + (3 \times 4)] + 20p[(3 \times 5) + (4 \times 5)] + (66pq - 58p)[(4 \times 6) + (4 \times 6)] = 3168pq - 1652p.$$

3) From the definition of first K hyper Banhatti index

$$HB_1(G) = \sum_{uv \in E(G)} [d_G(u) + d_G(v)]^2 = \sum_{ue \in E_{\{3,3\}}} [(d_G(u) + d_G(e))^2 + (d_G(v) + d_G(e))^2] + \sum_{ue \in E_{\{3,4\}}} [(d_G(u) + d_G(e))^2 + (d_G(v) + d_G(e))^2] + \sum_{ue \in E_{\{4,4\}}} [(d_G(u) + d_G(e))^2 + (d_G(v) + d_G(e))^2] = 12p[(3+4)^2 + (3+4)^2] + 20p[(3+5)^2 + (4+5)^2] + (66pq - 58p)[(4+6)^2 + (4+6)^2] = 13200pq - 6936p.$$

4) From the definition of second K hyper Banhatti index

$$HB_2(G) = \sum_{uv \in E(G)} [d_G(u)d_G(v)]^2 = \sum_{ue \in E_{\{3,3\}}} [(d_G(u)d_G(e))^2 + (d_G(v)d_G(e))^2] + \sum_{ue \in E_{\{3,4\}}} [(d_G(u)d_G(e))^2 + (d_G(v)d_G(e))^2] + \sum_{ue \in E_{\{4,4\}}} [(d_G(u)d_G(e))^2 + (d_G(v)d_G(e))^2] = 12p[(3 \times 4)^2 + (3 \times 4)^2] + 20p[(3 \times 5)^2 + (4 \times 5)^2] + (66pq - 58p)[(4 \times 6)^2 + (4 \times 6)^2] = 13200pq - 6936p.$$

$$(4 \times 5)^2] + (66pq - 58p)[(4 \times 6)^2 + (4 \times 6)^2] = 76\,032pq - 49\,132p.$$

Theorem 2 Let $G = L(F[p, q])$ be a line graph of V-Pantacenic nanotube. Then

$$BII_1(G) = 7^{24p} \times 8^{20p} \times 9^{20p} \times 10^{2(66pq - 58p)},$$

$$BII_2(G) = 12^{24p} \times (15)^{20p} \times (20)^{20p} \times (24)^{2(66pq - 58p)},$$

$$HBII_1(G) = 7^{48p} \times 8^{40p} \times 9^{40p} \times 10^{4(66pq - 58p)},$$

$$HBII_2(G) = 12^{48p} \times 15^{40p} \times 20^{40p} \times 24^{4(66pq - 58p)}.$$

Proof 1) From the definition of first multiplicative K Banahatti index

$$\begin{aligned} BII_1(G) &= \prod_{ue} [d_G(u) + d_G(e)] = \prod_{e=uv \in E_{[2,3]}} [d_G(u) + d_G(e)][d_G(v) + d_G(e)] \times \prod_{e=uv \in E_{[3,4]}} [d_G(u) + \\ &d_G(e)][d_G(v) + d_G(e)] \times \prod_{e=uv \in E_{[4,4]}} [d_G(u) + d_G(e)][d_G(v) + d_G(e)] = (3+4)^{12p} \times (3+4)^{12p} \times (3+5)^{20p} \times \\ &(4+5)^{20p} \times (4+6)^{(66pq-58p)} \times (4+6)^{(66pq-58p)} = 7^{24p} \times 8^{20p} \times 9^{20p} \times 10^{2(66pq-58p)}. \end{aligned}$$

2) From the definition of second multiplicative K Banahatti index

$$\begin{aligned} BII_2(G) &= \prod_{ue} d_G(u)d_G(e) = \prod_{e=uv \in E_{[2,3]}} [(d_G(u)d_G(e)) \times (d_G(v)d_G(e))] \times \prod_{e=uv \in E_{[3,4]}} [(d_G(u)d_G(e)) \times \\ &(d_G(v)d_G(e))] \times \prod_{e=uv \in E_{[4,4]}} [(d_G(u)d_G(e)) \times (d_G(v)d_G(e))] = (3 \times 4)^{12p} \times (3 \times 4)^{12p} \times (3 \times 5)^{20p} \times (4 \times \\ &5)^{20p} \times (4 \times 6)^{(66pq-58p)} \times (4 \times 6)^{(66pq-58p)} = 12^{24p} \times (15)^{20p} \times (20)^{20p} \times (24)^{2(66pq-58p)}. \end{aligned}$$

3) From the definition of first multiplicative K hyper Banahatti index

$$\begin{aligned} HBII_1(G) &= \prod_{ue} [d_G(u) + d_G(e)]^2 = \prod_{e=uv \in E_{[2,3]}} [d_G(u) + d_G(e)]^2 [d_G(v) + d_G(e)]^2 \times \prod_{e=uv \in E_{[3,4]}} [d_G(u) + \\ &d_G(e)]^2 [d_G(v) + d_G(e)]^2 \times \prod_{e=uv \in E_{[4,4]}} [d_G(u) + d_G(e)]^2 [d_G(v) + d_G(e)]^2 = [3+4]^2]^{12p} \times [(3+ \\ &4)^2]^{12p} \times [(3+5)^2]^{20p} \times [(4+5)^2]^{20p} \times [(4+6)^2]^{(66pq-58p)} \times [(4+6)^2]^{(66pq-58p)} = 7^{48p} \times 8^{40p} \times 9^{40p} \times \\ &10^{4(66pq-58p)}. \end{aligned}$$

4) From the definition of second multiplicative K hyper Banahatti index

$$\begin{aligned} HBII_2(G) &= \prod_{ue} (d_G(u)d_G(e))^2 = \prod_{e=uv \in E_{[2,3]}} [(d_G(u)d_G(e))^2 \times (d_G(v)d_G(e))^2] \times \prod_{e=uv \in E_{[3,4]}} [(d_G(u)d_G(e))^2 \times \\ &(d_G(v)d_G(e))^2] \times \prod_{e=uv \in E_{[4,4]}} [(d_G(u)d_G(e))^2 \times (d_G(v)d_G(e))^2] = [(3 \times 4)^2]^{12p} \times [(3 \times 4)^2]^{12p} \times [(3 \times 5)^2]^{20p} \times \\ &[(4 \times 5)^2]^{20p} \times [(4 \times 6)^2]^{(66pq-58p)} \times [(4 \times 6)^2]^{(66pq-58p)} = 12^{48p} \times 15^{40p} \times 20^{40p} \times 24^{4(66pq-58p)}. \end{aligned}$$

2 Conclusion

In this report we computed first K Banahatti index, second K Banahatti index, first K hyper Banahatti index, second K hyper Banahatti index, first multiplicative K Banahatti index, second multiplicative K Banahatti index, first multiplicative K hyper Banahatti index and second multiplicative K hyper Banahatti index of the line graph of V-Pantacenic nanotube. The following figure 3 and figure 4 show the dependence of first K Banahatti index and second K Banahatti index on the involved structural parameters p and q .




Figure 3 Plot of first K Banahatti index 3D(left), for $p=1$ (middle) and for $q=1$ (right)




Figure 4 Plot of second K Banahatti index 3D(left), for $p=1$ (middle) and for $q=1$ (right)

[参考文献]

- [1] WEST D B. Introduction to graph theory [M]. Upper Saddle River: Prentice hall, 2001.
- [2] RUCKER G, RUCKER C. On topological indices, boiling points, and cycloalkanes [J]. Journal of Chemical Information and Computer Sciences, 1999, 39 (5): 788–802.
- [3] KLAYZAR S, GUTMAN I. A comparison of the Schultz molecular topological index with the Wiener index [J]. Journal of Chemical Information and Computer Sciences, 1996, 36 (5): 1001–1003.
- [4] WIENER H. Structural determination of paraffin boiling points [J]. Journal of the American Chemical Society, 1947, 69 (1): 17–20.
- [5] DOBRYNIN A A, ENTRINGER R, GUTMAN I. Wiener index of trees: theory and applications [J]. Acta Applicandae Mathematica, 2001, 66 (3): 211–249.
- [6] GUTMAN I, POLANSKY O E. Mathematical concepts in organic chemistry [M]. Berlin: Springer Science & Business Media, 2012.
- [7] RANDIC M. Characterization of molecular branching [J]. Journal of the American Chemical Society, 1975, 97 (23): 6609–6615.
- [8] BOLLOBAS B, ERDOS P. Graphs of extremal weights [J]. Ars Combinatoria, 1998, 50: 225–233.
- [9] AMIC D, BESLO D, LUCIC B, et al. The vertex-connectivity index revisited [J]. Journal of Chemical Information and Computer Sciences, 1998, 38 (5): 819–822.
- [10] HU Y, LI X, SHI Y, et al. On molecular graphs with smallest and greatest zeroth-order general Randic index [J]. Match Communications in Mathematical and in Computer Chemistry, 2005, 54 (2): 425–434.
- [11] SARDAR M S, ZAFAR S, FARAHANI M R. The generalized Zagreb index of capra-designed planar benzenoid series [J]. Open Journal of Mathematical Sciences, 2017, 1 (1): 44–51.
- [12] REHMAN H M U, SARDAR R, RAZA A. Computing topological indices of hex board and its line graph [J]. Open Journal of Mathematical Sciences, 2017, 1 (1): 62–71.
- [13] SIDDIQUI H, FARAHANI M R. Forgotten polynomial and forgotten index of certain interconnection networks [J]. Open Journal of Mathematical Analysis, 2017 (1): 45–60.
- [14] SARDAR M S, PAN X F, GAO W, et al. Computing sanskruti index of titania nanotubes [J]. Open Journal of Mathematical Sciences, 2017, 1 (1): 126–131.
- [15] DE N. Hyper Zagreb index of bridge and chain graphs [J]. Open Journal of Mathematical Sciences, 2018, 2 (1): 1–17.
- [16] KULLI V R, CHALUVARAJU B, BOREGOWDA H S. Connectivity banhatti indices for certain families of benzenoid systems [J]. Journal of Ultra Chemistry, 2017, 13 (4): 81–87.
- [17] GAO W, MUZAFFAR B, NAZEER W. K-banhatti and K-hyper banhatti indices of dominating david derived network [J]. Open Journal of Mathematical Analysis, 2017 (1): 13–24.
- [18] KULLI V R. Multiplicative K hyper-banhatti indices and coindices of graphs [J]. International Journal of Mathematical Archive, 2016, 7 (6): 60–65.
- [19] GAO Y Y, NADEEM M F, ZAFAR S, et al. About the certain topological indices of the line graph of V-Pantacenic nanotube [J]. International Journal of Pharmaceutical Sciences and Research, 2017, 8 (12): 5349–5352.